

Z' Gauge Models from Strings*

Jose R. Espinosa^{a†}

^aDepartment of Physics, University of Pennsylvania
Philadelphia PA 19104, USA

Potentially realistic string models often contain additional abelian gauge factors besides the standard model group. The consequences of such extended gauge structure are manifold both for theory and phenomenology as I show focussing in the simplest case of just one additional non-anomalous $U(1)'$. First, I discuss the possible symmetry breaking patterns according to the scale at which the $U(1)'$ symmetry gets broken: in the first case, that scale must be below ~ 1 TeV to avoid fine-tuning; in the second case, the breaking can take place along a flat direction at an intermediate scale between the string scale and the electroweak scale. In both cases, I present a number of the generic implications expected, e.g. for the μ problem, Z' and Higgs physics, dark matter and fermion masses.

1. INTRODUCTION

Many string models predict a gauge group at the string scale which contains, besides the standard model group $G_{SM} = SU(3)_C \times SU(2)_L \times U(1)_Y$, extra $U(1)$ gauge factors. In this talk, I will examine the generic phenomenological and theoretical implications of this type of scenarios. For simplicity, I limit the analysis to one extra non-anomalous $U(1)'$. This is the simplest gauge extension of the SM one can consider and the phenomenology of particular Z' models has been studied extensively. Here, no attempt is made at building a concrete realistic string model but rather at exploring the generic implications of string inspired Z' models. As we shall see, such simple extended models can have manifold implications for, among others, the μ -problem, Higgs physics, exotic matter, fermion masses, dark matter, the spectrum of superpartners and cosmology. I will describe some of these implications in two different scenarios which can be distinguished by the scale of $U(1)'$ breaking. For more details the reader is directed to refs. [1–4].

Let me be more specific about the string input in the analysis. I will consider a class of

specific string vacua: orbifold models [5,6] with Wilson lines, in particular, models based on the free-fermionic construction (at special points of moduli space) [7–10]. I then assume:

- N=1 Supersymmetry
- The (visible sector) gauge group at $M_{string} (\simeq 5 \times 10^{17} \text{ GeV})$ is $G_{SM} \times U(1)'$, where the extra abelian factor has no anomaly.
- The matter content is that of the Minimal Supersymmetric Standard Model (MSSM) plus some number of SM singlets, S_i , which carry $U(1)'$ charges Q_i , plus some exotics (generally transforming non-trivially under G_{SM}). Both singlets and exotics appear generically in string models and can play an important role in breaking the $U(1)'$ symmetry.
- The gauge couplings g_3, g_2, g_1, g_1' (for $SU(3)_C, SU(2)_L, U(1)_Y$ and $U(1)'$ respectively, with proper normalization for g_1 and g_1' [2]) unify at M_{string} . This unification works only at the 10 – 15% level in the simplest models (in which the couplings meet at a scale below M_{string}). The small numerical inconsistency that results does not affect the generic predictions of these models that I will present.

*Talk given at 5th International Conference on Supersymmetries in Physics (SUSY 97), Philadelphia, PA, 27-31 May 1997

[†]Work supported by the U.S. Department of Energy Grant No. DOE-EY-76-02-3071.

- Supersymmetry breaking, the main uncertainty in extracting phenomenological consequences from string models, is parameterized by soft masses for scalars and gauginos and by trilinear masses associated to superpotential terms. Naturalness requires these masses (which I denote in general by m_{soft}) to be of the order of the electroweak scale.
- No bilinear mass terms are present in the superpotential and the Yukawa couplings at M_{string} are either zero or related to the unified gauge coupling g_0 by $\sqrt{2}$ factors (typically $\sim g_0\sqrt{2}, g_0$ or $g_0/\sqrt{2}$ in the fermionic $Z_2 \times Z_2$ orbifold constructions).

For fixed values of the parameters at the string scale one can address, by a renormalization group analysis (with the chosen parameters as boundary conditions) what are the low-energy implications typical of these scenarios.

2. RADIATIVE BREAKING OF $U(1)'$

To have an acceptable model, the extra $U(1)'$ gauge factor has to be broken at low energies. Generically, this breaking is achieved radiatively (in a way which parallels electroweak radiative breaking in the MSSM [11]) when the soft mass squared m_S^2 of some singlet S [charged under $U(1)'$] is driven to negative values in the infrared (an alternate breaking mechanism is discussed in the next section). This causes an instability of the potential for $\langle S \rangle = 0$, and S acquires a non-zero vacuum expectation value (VEV) thus breaking the $U(1)'$. The value of $\langle S \rangle$, and thus the breaking scale, is fixed by quartic D terms [from $U(1)'$] in the potential for the field S (alternatively there may be also quartic F -terms),

$$V(S) = m_S^2 S^2 + \frac{1}{2} g_1'^2 Q_S^2 S^4, \quad (1)$$

giving $g' \langle S \rangle \sim |m_S|$.

If the two Higgs doublets H_1, H_2 are charged under $U(1)'$ (the interesting case) electroweak breaking is influenced at tree-level by the breaking of $U(1)'$. In order to have a natural electroweak breaking scale (or equivalently, to avoid fine tuning the soft masses), the $U(1)'$ breaking scale cannot be much larger than $\mathcal{O}(1) \text{ TeV}$.

Thus, we see that the Z' models derived from strings naturally predict $M_{Z'} \lesssim \mathcal{O}(1) \text{ TeV}$ [1].

One exception to this generic result [besides the trivial one of having the $U(1)'$ decoupled from the electroweak Higgs sector] occurs if the $U(1)'$ is broken radiatively along a flat direction. For example, if the model contains two SM singlets $S_{1,2}$ with opposite $U(1)'$ charges, the field direction $S \equiv S_1 = S_2$ is D -flat and (assuming also F -flatness) the tree-level potential $V(S) = -m_S^2 S^2$ has no quartic coupling and is unstable. Eventually, the potential gets stabilized at very large values of the field S due to radiative corrections [the most important of which are encoded in the running mass $m_S^2(S)$] or by non-renormalizable operators (NROs) which can no longer be neglected for such large field values. In both cases the $U(1)'$ breaking scale is much larger than the electroweak scale (e.g. 10^{12} GeV). This is one example of an intermediate scale being generated in string models [12]. Still, these scenarios have also many implications for the low-energy effective theory. In the rest of the talk I will discuss separately the two general classes of scenarios just described: I) Z' below the TeV scale [2] and II) intermediate scale Z' [3]. For other related discussions in the literature see [13].

3. Z' BELOW THE TeV SCALE

Let us consider the simplest case, which requires the presence of just one singlet S to give a correct symmetry breaking pattern. We further assume that the $U(1)'$ charge assignments of S and the two Higgs doublets $H_{1,2}$ are such that the superpotential contains a term

$$W = h_s S H_1 \cdot H_2, \quad (2)$$

and thus $Q_S + Q_1 + Q_2 = 0$. After symmetry breaking, $\langle S \rangle = s/\sqrt{2}$, $\langle H_1^0 \rangle = v_1/\sqrt{2}$ and $\langle H_2^0 \rangle = v_2/\sqrt{2}$, the gauge group gets broken down to $U(1)_{em}$ with

$$M_W^2 = \frac{1}{4} g_2^2 (v_1^2 + v_2^2), \quad (3)$$

from which $v^2 \equiv v_1^2 + v_2^2 = (246 \text{ GeV})^2$, and

$$M_{Z-Z'}^2 = \begin{pmatrix} M_Z^2 & \Delta^2 \\ \Delta^2 & M_{Z'}^2 \end{pmatrix}, \quad (4)$$

where

$$M_Z^2 = \frac{1}{4}G^2(v_1^2 + v_2^2), \quad (5)$$

$$M_{Z'}^2 = g_1'^2(v_1^2 Q_1^2 + v_2^2 Q_2^2 + s^2 Q_S^2), \quad (6)$$

$$\Delta^2 = \frac{1}{2}g_1' G(v_1^2 Q_1 - v_2^2 Q_2). \quad (7)$$

Here we use $G^2 = g_2^2 + g_Y^2 = g_2^2 + 3g_1^2/5$.

From Eqs. (4) and (7) we see that it is possible to accomodate a very small $Z - Z'$ mixing angle (the experimental upper bound is $\mathcal{O}(10^{-3})$ in typical models [14]) due to a cancellation in the off-diagonal term Δ^2 for a particular value of $\tan\beta \equiv v_2/v_1 \sim \sqrt{|Q_1/Q_2|}$. The presence of the singlet VEV s , on the other hand, helps in giving to Z' a sufficiently large mass to evade collider constraints [19]. In ref. [2] two phenomenologically viable scenarios were identified:

- **(a) A -driven scenario.** Symmetry breaking is driven by a large value of the trilinear soft mass associated to the superpotential term (2) [this is similar to $SU(3)_C$ color breaking in the MSSM for large values of A_Q , the trilinear coupling associated to the quark Yukawa couplings], that is, $A \gg M_{1/2}, m_0$, where $M_{1/2}$ represents a typical gaugino mass and m_0 a scalar soft mass. In this scenario, the minimum of the potential has $v_1 \simeq v_2 \simeq s$. This is shown in figure 1, where the adimensional ratio $f_i = v_i/m_0$ is plotted as a function of $c_A = A/m_0$ in some particular example. For $Q_1 = Q_2$, values of $\tan\beta$ close to 1 would give a small enough mixing angle.
- **(b) Large s scenario.** ($s \gg v_1, v_2$) This case corresponds to a hierarchy in the masses $M_{Z'} \gg M_Z$. The electroweak scale, as represented by M_Z , is light due to accidental cancellations among soft masses, which are typically of the order $M_{Z'}$. To avoid fine-tuning, $M_{Z'}$ cannot be much larger than $\mathcal{O}(1 \text{ TeV})$ as we already discussed before.

3.1. Soft Masses at M_{string}

One can then proceed to determine what pattern of soft supersymmetry breaking terms at the




Figure 1. c_A dependence of the dimensionless field VEVs for the A -driven scenario with $h_s = 0.7$ and $Q_1 = Q_2 = -1$.

string scale will lead, via RG evolution, to the two low-energy acceptable scenarios described previously.

With the minimal particle content (MSSM + singlet) it is not possible to achieve the correct symmetry breaking if universal boundary conditions for the soft masses at M_{string} are used. Both scenarios, (a) and (b), can however be reached with non-universal boundary conditions. The allowed parameter space is somewhat constrained and typically corresponds to gaugino masses much smaller than the scalar masses.

It is easier to have an acceptable symmetry breaking pattern in the presence of exotics, in particular if the singlet S has couplings to them of the form

$$W = h_E S E_1 E_2. \quad (8)$$

The large value of the Yukawa coupling h_E ($\sim$

g_0) triggers easily $U(1)'$ breaking and the large s scenario [(b)] can be reached even with universal soft masses. The large A scenario [(a)] is more constrained and some degree of nonuniversality is required.

3.2. The μ -problem

In the presence of the superpotential term $h_s S H_1 \cdot H_2$, the μ term of the MSSM Higgs potential is generated dynamically by the VEV of the singlet S as

$$\mu_{eff} = h_s \langle S \rangle. \quad (9)$$

As discussed before, $\langle S \rangle$ lies at the electroweak scale, so that μ_{eff} is of the correct order of magnitude to give a proper electroweak breaking [the same mechanism generates a B term $\sim h_s \langle S \rangle A \sim m_{soft}$]. This solves the μ problem [15] of the MSSM (where μ is a SUSY parameter which does not know about the electroweak scale and has to be set of that order by hand).

This solution to the μ problem is similar in spirit to the mechanism in the Next to Minimal Supersymmetric Standard Model (NMSSM) [16] in which the MSSM is extended by a chiral singlet but no additional $U(1)$ gauge symmetry is present. The advantage of having a $U(1)'$ present is that there is no cosmological problem with the formation of domain walls after symmetry breaking: the discrete Z_3 symmetry of the NMSSM responsible for the domains is embedded in $U(1)'$.

This solution is complementary to the Giudice-Masiero mechanism [17], in the sense that the $U(1)'$ symmetry would forbid a $H_1 H_2$ term in the Kähler potential, which is in the basis of that mechanism.

The two scenarios (a) and (b) would correspond to a small and large μ_{eff} respectively. In the first scenario (A dominated) light charginos and neutralinos are expected, with obvious implications for LEP II searches [20].

3.3. Higgs Physics

The (CP conserving) Higgs sector contains two doublets H_1, H_2 and one singlet S making up a total of 10 degrees of freedom. After spontaneous symmetry breaking, 4 d.o.f. are eaten by the longitudinal components of Z, Z' and $W^\pm$. The 6

d.o.f. left appear as physical Higgs bosons: three (CP even) scalars H_i^0 , one (CP odd) pseudoscalar A^0 and one charged pair $H^\pm$. That is, the model contains just one scalar more than in the MSSM.

The pattern of Higgs masses is very different in the two scenarios (a) and (b):

In the A -dominated scenario, all Higgs masses are controlled by one single mass scale v and three adimensional parameters: h_s, g'_1 and Q_1 . The spectrum is therefore light, although all Higgses can still evade detection at LEP II. One typical example, for $h_s = 0.7, g'_1{}^2 = (5/3)G^2 \sin^2 \theta_W$ and $Q_1 = -1$, has $m_{H^\pm} = 146 \text{ GeV}$, $m_{A^0} = 211 \text{ GeV}$, $m_{H_3^0} = 230 \text{ GeV}$, $m_{H_2^0} = 152 \text{ GeV}$ and $m_{H_1^0} = 122 \text{ GeV}$ (tree-level masses).

In the large s scenario there are three mass scales in the Higgs spectrum. Typically $m_{H_3^0} \simeq M_{Z'}$ (with H_3^0 singlet dominated), $m_{A^0} \gg M_Z$, with $(H_2^0, A^0; H^\pm)$ forming a nearly degenerate doublet decoupled from electroweak symmetry breaking, and H_1^0 remains at the electroweak scale.

An upper bound on the tree-level mass of H_1^0 can be easily obtained as

$$m_{H_1^0}^2 \leq M_Z^2 \cos^2 2\beta + \frac{1}{2} h_s^2 v^2 \sin^2 2\beta + g'_1{}^2 \overline{Q}_H^2 v^2, \quad (10)$$

with $\overline{Q}_H = Q_1 \cos^2 \beta + Q_2 \sin^2 \beta$. The first term in Eq. (10), coming from $SU(2)_L \times U(1)_Y$ D -terms, is the MSSM piece. The second, from F -terms, appears also in the NMSSM but the third piece, from $U(1)'$ D -terms, is a particular feature of this type of models [21–23]. More refined bounds can be obtained under extra assumptions for the parameters of the model (see [2]). This tree-level bound can be very large; 170 GeV is a typical example (it is usually forgotten that the bound $\sim 150 \text{ GeV}$ for any perturbative SUSY model [23] applies only if the gauge sector is not extended) so that H_1^0 can easily evade detection at LEP II.

3.4. Superpartner Spectrum

The spectrum of superpartners is also typically light for the A -dominated scenario (with all particles expected to lie at the electroweak breaking scale) while in the large s scenario soft masses are generically larger, giving a decoupled SUSY mass

pattern.

There is, however, an extra effect associated with $U(1)'$ D -term contributions to the masses. The squared mass of any scalar ϕ_i (with $U(1)'$ charge Q_i) is shifted by the amount

$$\delta m_i^2 = \frac{1}{2} g_1'^2 Q_i (Q_1 v_1^2 + Q_2 v_2^2 + Q_S s^2). \quad (11)$$

This can be a sizable effect that can cause some superpartner to be lighter. In general, the mass sum-rules that can be derived in particular models will be violated by these contributions, leaving a characteristic imprint in the mass spectrum [24].

3.5. Dark Matter

The neutralino sector in these models is composed of $\tilde{B}$, $\tilde{W}_3$, $\tilde{H}_1^0$, $\tilde{H}_2^0$ plus $\tilde{B}'$ and $\tilde{S}$. In most of the parameter space the LSP (which is the natural candidate for dark matter when R -parity is conserved) is the lightest neutralino with composition dominated by $\tilde{B}$. However, in some regions of parameter space (with large s and large gaugino masses) the lightest neutralino is $\tilde{S}$ -dominated making an atypical dark matter candidate [25] with interesting rates for direct detection ($\sim 10^{-2}$ events/kg/day in ^{73}Ge detectors).

4. INTERMEDIATE SCALE Z'

In this case, at least two SM singlets, S_1, S_2 , with $Q_1 Q_2 < 0$ are required. Calling S the flat direction determined by the condition $|Q_1| S_1^2 = |Q_2| S_2^2$ (we assume also F -flatness along S) the renormalization-group improved potential for S is simply

$$V(S) = m_S^2(S) S^2. \quad (12)$$

If the couplings of S are such that $m_S^2 < 0$ in the infrared (we assume $m_S^2 > 0$ at the string scale) the shape of the potential is as depicted in figure 2 (solid line labeled RAD). The potential tracks the running of m_S^2 (dashed line) and is stabilized when m_S^2 turns positive at the scale μ_{RAD} . The minimum of the potential occurs at some intermediate scale $m_I \lesssim \mu_{RAD} \gg M_Z$.




Figure 2. Effective potential (solid lines) along the flat direction: **1.** (marked *RAD*) in the case of stabilization by the running of m_S^2 (dashed line) and **2.** (marked *NRO*) in the case of stabilization by non-renormalizable operators below the scale μ_{RAD} .

If μ_{RAD} is very large, non-renormalizable operators like

$$W_{NRO} \sim \alpha_K \left(\frac{S}{M} \right)^K S^3, \quad (13)$$

which are generically present after integrating out physics at the string scale ($M \simeq M_{string}$) cannot be neglected in the determination of the minimum. In (13), we write the NRO term, originally given by powers of S_1 and/or S_2 , projected along the flat direction S . These terms contribute to the potential a positive-definite term

$$\delta V(S) \sim \left(\frac{S^{2+K}}{M^K} \right)^2, \quad (14)$$

that lifts the flat direction and can stabilize the potential below μ_{RAD} . The shape of the potential

in this case is represented in figure 2 by the solid line labeled NRO.

We can distinguish then two different classes of models according to the mechanism for stabilization of the potential:

- **1.** Stabilization by pure running of m_S^2 (typical case for $\mu_{RAD} \lesssim 10^{12} \text{ GeV}$). The minimum is fixed at $\langle S \rangle \sim \mu_{RAD}$.
- **2.** Stabilization by NROs (typical case for $\mu_{RAD} \gtrsim 10^{12} \text{ GeV}$). The minimum is fixed at

$$\langle S \rangle \sim \mu_K \equiv (m_{soft} M^K)^{\frac{1}{K+1}} < \mu_{RAD}. \quad (15)$$

4.1. Radiative Breaking

Both scenarios, 1 and 2, are based on radiative breaking of the $U(1)'$ symmetry along the flat direction S . The precise condition that has to be satisfied at the electroweak scale to ensure the breaking (the $m_S^2 < 0$ condition of the previous subsection) is

$$\frac{m_1^2}{|Q_1|} + \frac{m_2^2}{|Q_2|} < 0, \quad (16)$$

where m_1^2, m_2^2 are the (electroweak scale) soft masses of S_1 and S_2 respectively. From Eq. (16) we see that $m_1^2 < 0$ and/or $m_2^2 < 0$ are required [and a hierarchy in the charges can make (16) easier to fulfill].

This breaking can be easily achieved when S_1 (and/or S_2) couples to exotics in a superpotential term

$$W = h_E S_i E_1 E_2. \quad (17)$$

In [3] different cases (with different quantum numbers and multiplicities for the exotic fields E_i) were studied. It was found that a correct breaking can be obtained even in the case of universal soft masses at the string scale (provided the multiplicity of the exotics is large enough). By varying the parameters of the model the scale μ_{RAD} can be moved in the wide range 10^5 GeV to 10^{16} GeV .

4.2. Low-Energy Effective Theory

In both cases, 1 and 2, the $U(1)'$ breaking takes place at an intermediate scale (e.g. 10^{12} GeV)

much larger than the electroweak scale. The Z' boson (and the exotics coupled to the flat direction) is thus very heavy (this can have interesting effects for gauge coupling unification [26]). The effective theory at the TeV scale is just the MSSM, but with an extra chiral multiplet $\hat{S}$, remnant of $U(1)'$ breaking. However, the interactions of $\hat{S}$ with MSSM fields are suppressed by powers of the intermediate scale. Nevertheless, these scenarios are interesting because many parameters of the MSSM can be determined by intermediate scale physics.

For example, the mass spectrum of superpartners is affected by the intermediate scale $U(1)'$ breaking. At the renormalizable level, the only interaction between the MSSM fields and the intermediate scale fields arises from the $U(1)'$ D -terms in the scalar potential. The resulting effect after integrating out the fields which have heavy intermediate scale masses [24] is a shift of the soft masses of MSSM fields charged under the extra $U(1)'$:

$$\delta m_i^2 = -Q_i \frac{m_1^2 - m_2^2}{Q_1 - Q_2}. \quad (18)$$

4.3. The μ parameter

In these models, the μ parameter can be generated dynamically by the VEV of S through non-renormalizable operators in the superpotential that couple the MSSM Higgs doublets to the flat direction [15]:

$$W = h S H_1 \cdot H_2 \left(\frac{S}{M} \right)^P. \quad (19)$$

From this term, the following effective μ parameter results

$$\mu_{eff} = h \frac{\langle S \rangle^{P+1}}{M^P}. \quad (20)$$

Its order of magnitude in both scenarios, 1 and 2, is:

- **1. Radiative Stabilization.** In this case $\langle S \rangle \sim \mu_{RAD}$. Depending on the value of P in Eq. (20), the value of μ_{RAD} must be tuned to give $\mu_{eff} \sim m_{soft}$:

$$\mu_{RAD} \sim (m_{soft} M^P)^{\frac{1}{P+1}}. \quad (21)$$

- **2. Stabilization by NROs.** Eq. (21) is exactly of the form expected for μ_K [see Eq. (15)], the scale of the minimum in these scenarios. In other words, for K in Eq. (13) equal to P in Eq. (19), one gets automatically $\mu_{eff} \sim m_{soft}$. The same applies to the B -term in the MSSM Higgs potential, which can be generated by $\langle S \rangle$ in a similar way.

4.4. Non-Renormalizable Operators

As we have seen, NROs play a crucial role in these intermediate scale models. Interestingly, NROs are calculable in a large class of string models [27,28]. Writing

$$W = \alpha_K \left(\frac{S}{M_{Pl}} \right)^K \equiv \left(\frac{S}{M_K} \right)^K, \quad (22)$$

the coefficient α_K is basically given by a complicated world-sheet integral and some Clebsch-Gordan coefficients. Reabsorbing α_K in the mass M_K , the general tendency is that M_K increases with K so that the weight of higher order terms is further suppressed [27,28,3].

It is also important to determine which NROs will be present, that is, what values of the powers P (for the μ parameter) and K (to fix the minimum) will be allowed. The values of these powers are constrained by

- $U(1)'$ gauge invariance. For example, if $Q_1 = 4/5$ and $Q_2 = -1/5$,

$$W \sim (S_1 S_2^4)^n \sim S^{5n}, \quad (23)$$

so that $K = 5n$.

- World-sheet selection rules, e.g. in free-fermionic models, can forbid terms otherwise consistent with all gauge symmetries (see, for example, [28,29]). So, one should keep in mind that not all the NROs satisfying all gauge symmetry constraints will be necessarily present in the superpotential. This is a typical stringy feature which is in contrast with the field theory lore.

4.5. Fermion Masses

The large value of string Yukawa couplings offers a natural explanation for the large value of the top mass [18]. The light fermions would have no Yukawa couplings in the superpotential and would therefore be massless, which is a good first order approximation. Models with an intermediate scale provide a mechanism to give small non-zero masses to these fermions using NROs like

$$\begin{aligned} W_{u_i} &\sim H_2 \cdot Q_i U_i^c \left(\frac{S}{M} \right)^{P'_{u_i}}, \\ W_{d_i} &\sim H_1 \cdot Q_i D_i^c \left(\frac{S}{M} \right)^{P'_{d_i}}, \\ W_{e_i} &\sim H_1 \cdot L_i E_i^c \left(\frac{S}{M} \right)^{P'_{e_i}}, \end{aligned} \quad (24)$$

where i is a generation index and I use standard notation for the MSSM chiral supermultiplets. Small Yukawa couplings are generated after $U(1)'$ symmetry breaking:

$$Y \sim \left(\frac{\langle S \rangle}{M} \right)^{P'}. \quad (25)$$

Limiting the discussion to scenarios with NRO stabilization, which give more definite predictions, it was shown in [3] how the family pattern of quark and lepton masses [30] can be qualitatively reproduced for particular choices of the powers P' in Eq. (25) and K in Eq. (13). In particular, as shown in Table 1, $K = 5, 6$ in combination with $P' = 1$ for the second family and $P' = 2$ for the first are satisfactorily close to the observed pattern. The mass of the light fermions of the third family do not fit quite as well and call for some other mechanism. However, given the roughness of the estimates and the simplicity of the model, the overall pattern of masses is quite encouraging.

There are also interesting implications for neutrino masses [31]. A very light (non-seesaw) doublet neutrino Majorana mass is possible from the superpotential

$$W \sim \frac{H_2 \cdot L_i}{M} \left(\frac{S}{M} \right)^{P''_{L_i L_i}}, \quad (26)$$

Table 1

Non-Renormalizable MSSM mass terms via $\langle S \rangle$. For $m_{soft} \sim 100$ GeV, $M \sim 3 \times 10^{17}$ GeV.

	P or P'	$K = 2$	$K = 3$	$K = 4$	$K = 5$	$K = 6$	$K = 7$
$\left(\frac{m_{soft}}{M}\right)^{\frac{1}{K+1}}$		7×10^{-6}	1×10^{-4}	8×10^{-4}	3×10^{-3}	6×10^{-3}	1×10^{-2}
$\langle S \rangle$ (GeV)		2×10^{12}	4×10^{13}	2×10^{14}	8×10^{14}	2×10^{15}	3×10^{15}
$\frac{\mu_{eff}}{m_{soft}}$	$K - 1$	1×10^5	7×10^3	1×10^3	400	200	90
	K	1	1	1	1	1	1
	$K + 1$	7×10^{-6}	1×10^{-4}	8×10^{-4}	3×10^{-3}	6×10^{-3}	1×10^{-2}
$\frac{m_{Q,L}}{\langle H_i \rangle}$	0	1	1	1	1	1	1
	1	7×10^{-6}	1×10^{-4}	8×10^{-4}	3×10^{-3}	6×10^{-3}	1×10^{-2}
	2	5×10^{-11}	2×10^{-8}	6×10^{-7}	7×10^{-6}	4×10^{-5}	1×10^{-4}
	3	3×10^{-16}	2×10^{-12}	5×10^{-10}	2×10^{-8}	2×10^{-7}	2×10^{-6}
	4	2×10^{-21}	3×10^{-16}	4×10^{-13}	5×10^{-11}	1×10^{-9}	2×10^{-8}
	5	2×10^{-26}	5×10^{-20}	3×10^{-16}	1×10^{-13}	9×10^{-12}	2×10^{-10}

but the generated mass is too small ($< 10^{-4}$ eV) to be relevant for dark matter or MSW conversions in the Sun. If right-handed neutrino chiral multiplets, ν_i^c , are introduced in the model, then Majorana and Dirac neutrino mass terms can be generated via

$$W_D \sim H_2 \cdot L_i \nu_i^c \left(\frac{S}{M} \right)^{P'_{L_i \nu_i^c}},$$

$$W_M \sim \nu_i^c \nu_i^c S \left(\frac{S}{M} \right)^{\bar{P}_{\nu_i^c \nu_i^c}}. \quad (27)$$

The superpotential W_D can yield Dirac neutrino masses heavier than those coming from Eq. (26). For example, taking $K = 5$ and $P'_{L_i \nu_i^c} = 4$ or 5 gives masses $m_{L_i \nu_i^c} = 0.9$ eV or 10^{-2} eV, respectively, which are in the interesting range for solar and atmospheric neutrinos, oscillation experiments and dark matter.

The superpotential W_M , on the other hand, can give a Majorana mass to the singlet neutrinos ν_i^c which can be very large or small, depending on the sign of $\bar{P}_{\nu_i^c \nu_i^c} - K$. Combining the effects of W_M and W_D , light neutrinos can be produced by the standard seesaw mechanism, with mass

$$m_{seesaw}^{light} \sim m_{L_i \nu_i^c}^2 / m_{\nu_i^c \nu_i^c}$$

$$\sim m_{soft} \left(\frac{m_{soft}}{M} \right)^{\frac{2P'_{L_i \nu_i^c} + K - \bar{P}_{\nu_i^c \nu_i^c}}{K+1}}. \quad (28)$$

With the choice $K = 5$ and $P'_{L_i \nu_i^c} = P'_i = \{2, 1\}$ for $i = 1, 2$, respectively and with either $P'_{L_3 \nu_3^c} = 1$ or $P'_{L_3 \nu^c} = P'_{u_3} = 0$ (involving a renormalizable Dirac neutrino term) the light eigenvalues of the three generations fall into the hierarchy of 3×10^{-5} eV, 1×10^{-2} eV, and either 1×10^{-2} eV or 5 eV for $\bar{P}_{\nu_i^c \nu_i^c} = P'_{L_i \nu_i^c} + 1$. This range is again of interest for laboratory and non-accelerator experiments.

5. SUMMARY AND CONCLUSIONS

A large class of potentially realistic string models predict a gauge group of the observable sector with additional abelian factors besides the standard group. In this talk I have discussed some phenomenological and theoretical implications of this type of models concentrating for simplicity in the simplest case, with just one additional (non-anomalous) $U(1)'$. This modest extension already has a plethora of interesting consequences, some of which I have described. The goal has been to explore the generic features of these models rather than to construct a realistic string model. In this respect one can divide the scenarios in two different classes according to the scale at which the $U(1)'$ symmetry gets broken.

Generically, $U(1)'$ breaking takes place below the TeV scale much in the same way that electroweak breaking occurs. Actually, when the two MSSM Higgs doublets that break $SU(2)_L \times U(1)_Y$

are charged under $U(1)'$ (which is the case of interest) the breaking of $U(1)'$ is linked to electroweak breaking. To avoid fine-tuning the parameters of the model, the scale of $U(1)'$ breaking cannot be much larger than the Fermi scale. Thus, the new Z' boson is expected to be on the reach of the next generation of colliders. An acceptable symmetry breaking pattern requires the presence of at least one standard model singlet S charged under the $U(1)'$ and taking a VEV. If the coupling $SH_1 \cdot H_2$ is present in the superpotential, the μ parameter is generated dynamically and with the correct order of magnitude.

In this setting, two main possibilities exist for a phenomenologically viable scenario: a) symmetry breaking driven by the trilinear soft term associated to $SH_1 \cdot H_2$ and b) radiative breaking with a singlet VEV large compared to the electroweak scale (by say, one order of magnitude). Both cases can give an acceptably small $Z - Z'$ mixing angle and evade collider bounds on $M_{Z'}$. Case a) corresponds to a relatively light Z' while case b) gives a heavy Z' . With minimal particle content (no exotics), both scenarios can arise as the low energy limit of a string model, if some degree of non-universality in the soft masses is present at the string scale. In the presence of exotics, the large s scenario can be obtained even keeping universal boundary conditions. Besides solving the μ problem, these scenarios have interesting implications for Higgs physics, the mass spectrum of superpartners and dark matter, among others.

An exception to the generic case described above can occur if the $U(1)'$ is broken along a flat direction. Such flat directions exist if at least two SM singlets have $U(1)'$ charges of opposite signs (F -flatness is also required). In that case, the field along the flat direction, S , can take a very large VEV. This is what happens if the potential $V(S)$ gets destabilized at the origin by a negative soft mass squared m_S^2 for S in the infrared. Two main classes of scenarios are possible depending on the mechanism that stabilizes the potential at large S and fixes the VEV [and thus the scale of $U(1)'$ breaking]: in one, the potential is stabilized by m_S^2 running to positive values in the ultraviolet; in the second, stabilization is due to nonrenormalizable operators. This latter case is more pre-

dictive as the scale involved is fixed mainly by the (discrete) order of the NROs. In both cases the low-energy effective theory is simply the MSSM but many of its parameters are influenced by the intermediate scale breaking of the $U(1)'$. There are, for example, effects on the soft masses for superpartners, which receive new $U(1)'$ contributions. In addition, NROs coupling MSSM fields to the flat direction can offer an explanation for the origin of the μ parameter, as well as for the hierarchy of the observed fermion masses (and there are also interesting implications for neutrino masses).

In conclusion, it is worth pursuing the detailed study of such extensions of the minimal supersymmetric standard model. These simple models are not only well motivated from the string perspective but have a remarkably large number of potentially interesting implications, both phenomenologically and theoretically.

Acknowledgments

I thank G. Cleaver, M. Cvetič, D.A. Demir, B. de Carlos, L. Everett and P. Langacker for very enjoyable collaborations on the topics covered.

REFERENCES

1. M. Cvetič and P. Langacker, Phys. Rev. **D54**, 3570 (1996), and Mod. Phys. Lett. **11A**, 1247 (1996).
2. M. Cvetič, D.A. Demir, J.R. Espinosa, L. Everett and P. Langacker, Preprint UPR-0737-T [hep-ph/9703317] to appear in Phys. Rev. **D**.
3. G. Cleaver, M. Cvetič, J.R. Espinosa, L. Everett and P. Langacker, Preprint UPR-0750-T [hep-ph/9705391].
4. M. Cvetič and P. Langacker, Preprint UPR-0761-T, [hep-ph/9707451].
5. L. Dixon, J. A. Harvey, C. Vafa and E. Witten, Nucl. Phys. **B261**, 678 (1985); **B274**, 285 (1986).
6. A. Font, L. E. Ibáñez, F. Quevedo and A. Sierra, Nucl. Phys. **B331**, 421 (1991).
7. I. Antoniadis, C. Bachas and C. Kounnas, Nucl. Phys. **B289**, 87 (1987); H. Kawai, D. Lewellen and S.H.-H. Tye, Phys. Rev.

- Lett. **57**, 1832 (1986); Phys. Rev. **D34**, 3794 (1986).
8. I. Antoniadis, J. Ellis, J. Hagelin and D. Nanopoulos, Phys. Lett. **B231**, 65 (1989).
 9. A. Faraggi, D.V. Nanopoulos and K. Yuan, Nucl. Phys. **B335**, 347 (1990); A. Faraggi, Phys Lett. **B278**, 131 (1992).
 10. S. Chaudhuri, S.-W. Chung, G. Hockney and J. Lykken, Nucl. Phys. **B456**, 89 (1995); S. Chaudhuri, G. Hockney and J. Lykken, Nucl. Phys. **B469**, 357 (1996).
 11. L. Ibáñez and C. López, Nucl. Phys. **B233**, 511 (1984); L. Ibáñez, C. López and C. Muñoz, Nucl. Phys. **B256**, 218 (1985).
 12. M. Dine, V. Kaplunovsky, M. Mangano, C. Nappi and N. Seiberg, Nucl. Phys. **B259**, 549 (1985); G. Costa, F. Feruglio, F. Gabiani and F. Zwirner, Nucl. Phys. **B286**, 325 (1987); J. Ellis, K. Enqvist, D.V. Nanopoulos and K. Olive, Phys. Lett. **188B**, 415 (1987); R. Arnowit and P. Nath, Phys. Rev. Lett. **60**, 1817 (1988).
 13. E. Keith and E. Ma, [hep-ph/9704441]; D. Suematsu and Y. Yamagishi, Int. J. Mod. Phys. **A10**, 4521 (1995); V. Jain and R. Shrock, Phys. Lett. **B352**, 83 (1995) and ITP-SB-95-22, [hep-ph/9507238]; Y. Nir, Phys. Lett. **354**, 107 (1995); J.D. Lykken, Preprint FERMILAB-CONF-96-344-T, [hep-ph/9610218]; J.F. Gunion, L. Roszkowski and H.E. Haber, Phys. Lett. **189B**, 409 (1987).
 14. See the Appendix of M. Cvetič and P. Langacker, Mod. Phys. Lett. **11A**, 1247 (1996), and references therein.
 15. J. E. Kim and H. P. Nilles, Phys. Lett. **B138**, 150 (1984).
 16. P. Fayet, Nucl. Phys. **B90**, 104 (1975); H.-P. Nilles, M. Srednicki and D. Wyler, Phys. Lett. **B120**, 346 (1983); J.-M. Frère, D.R.T. Jones and S. Raby, Nucl. Phys. **B222**, 11 (1983); J.-P. Derendinger and C.A. Savoy, Nucl. Phys. **B237**, 307 (1984); L. Durand and J.L. López, Phys. Lett. **B217**, 463 (1989); M. Drees, Int. J. Mod. Phys. **A4**, 3645 (1989); J. Ellis et al., Phys. Rev. **D39**, 844 (1989); S. F. King and P. L. White, Phys. Rev. **D52**, 4183 (1995); U. Ellwanger, M. Rausch de Traubenberg and C.A. Savoy, Nucl. Phys. **B492**, 21 (1997).
 17. G. F. Giudice and A. Masiero, Phys. Lett. **B206**, 480 (1988). See also J. A. Casas and C. Muñoz, Phys. Lett. **B306**, 288 (1993).
 18. See e.g., A.E. Faraggi, Phys.Lett. **B377**, 43(1996).
 19. K. Maeshima, Proceedings of the 28th International Conference on High Energy Physics (ICHEP'96), Warsaw, Poland, July 25-31, 1996.
 20. Special CERN particle physics seminar on physics results from the LEP run at $\sqrt{s} = 172$ GeV (25 Feb. 1997) by the four LEP collaborations.
 21. H.E. Haber and M. Sher, Phys. Rev. **D35** 2206 (1987); M. Drees, Phys. Rev. **D35**, 2910 (1987).
 22. D. Comelli and C. Verzegnassi, Phys. Rev. **D47**, 764 (1993) and Phys. Lett. **B303**, 277 (1993).
 23. J.R. Espinosa and M. Quirós, Phys. Lett. **B279**, 92 (1992) and Phys.Lett. **B302**, 51 (1993); G. Kane, C. Kolda, J.D. Wells, Phys. Rev. Lett. **70**, 2686 (1993); D. Comelli and J.R. Espinosa, Phys. Lett. **B388**, 793 (1996).
 24. M. Drees, Phys. Lett. **B181**, 279 (1986); H.-C. Cheng and L. Hall, Phys. Rev. **D51**, 1337 (1995); C. Kolda and S. Martin, Phys. Rev. **D53**, 3871 (1996).
 25. B. de Carlos and J.R. Espinosa, Preprint UPR-0749-T [hep-ph/9705315] to appear in Phys. Lett. **B**.
 26. See K.R. Dienes and A. Faraggi, Nucl. Phys. **B457**, 409 (1995), and references therein; P. Binetrui and P. Langacker, in preparation.
 27. S. Kalara, J. López, and D. Nanopoulos, Nucl. Phys. **B353**, 650 (1991).
 28. A. Faraggi, Nucl. Phys. **B487**, 55 (1997).
 29. J. Rizos and K. Tamvakis, Phys. Lett. **B262**, 227 (1991).
 30. R. Barnett *et al.*, *Reviews of Particle Properties*, Phys. Rev. **D54**, 1 (1996).
 31. For detailed reviews, see G. Gelmini and E. Roulet, Rept. Prog. Phys. **58**, 1207 (1995); P. Langacker in *Testing The Standard Model*, ed. M. Cvetič and P. Langacker (World, Singapore, 1991) p. 863.